

February 1, 2008

On quantum algebra symmetries of discrete Schrödinger equations

A. Ballesteros[†], F.J. Herranz[†], J. Negro[‡] and L.M. Nieto[‡]

[†] *Departamento de Física*
Universidad de Burgos, E-09001 Burgos, Spain

[‡] *Departamento de Física Teórica*
Universidad de Valladolid, E-47011 Valladolid, Spain

Abstract

Two non-standard quantum deformations of the (1+1) Schrödinger algebra are identified with the symmetry algebras of either a space or time uniform lattice discretization of the Schrödinger equation. For both cases, the deformation parameter of the corresponding Hopf algebra can be interpreted as the step of the lattice. In this context, the introduction of nonlinear maps defining Schrödinger and $sl(2, \mathbb{R})$ quantum algebras with classical commutation rules turns out to be relevant. The problem of finding a quantum algebra linked to the full space-time discretization is also discussed.

PACS numbers: 02.20.Sv, 03.65.Fd

Keywords: quantum algebras, Lie symmetries, Schrödinger equations, space-time lattices, uniform grids

I. INTRODUCTION

Since they were introduced, quantum algebras have been connected with different versions of space-time lattices through several algebraic constructions that have no direct relationship with the usual Lie symmetry theory.¹⁻³ Recent works^{4,5} have also developed new techniques for dealing with the symmetries of difference or differential-difference equations and have tried to adapt in this field the standard methods that have been so successful with differential equations. An exhaustive study for the discretization on q -lattices of classical linear differential equations has shown that their symmetries obeyed to q -deformed commutation relations with respect to the Lie algebra structure of the continuous symmetries.⁶⁻⁸ However, Hopf algebra structures underlying these q -symmetry algebras have been not found.

When the discretization of linear equations is made on uniform lattices it is well known that the relevant symmetries preserve the Lie algebra structure.^{9,10} Perhaps, this is the reason why the symmetry approach to these equations has never been related to quantum algebras. We will address this question in this paper for the discrete (1+1)-Schrödinger equation. In this case we will show that Lie algebras of (discrete) symmetries can be put in correspondence to non-trivial quantum Hopf algebra structures of the non-standard (or triangular) type.¹¹⁻¹⁶

To begin with, let us consider the following discrete version of the heat or (time imaginary) Schrödinger equation (SE) on a two-dimensional uniform lattice⁹

$$(\Delta_x^2 - 2m\Delta_t)\phi(x, t) = 0. \quad (1)$$

The difference operators Δ_x and Δ_t which appear in (1) can be expressed in terms of shift operators $T_x = e^{\sigma\partial_x}$ and $T_t = e^{\tau\partial_t}$ as

$$\Delta_x = \frac{T_x - 1}{\sigma} \quad \Delta_t = \frac{T_t - 1}{\tau}, \quad (2)$$

where the parameters σ and τ are the lattice constants in the space x and time t directions, respectively. The action of Δ_x or Δ_t on a function $\phi(x, t)$ consists in a discrete derivative, which in the limit $\sigma \rightarrow 0$ and $\tau \rightarrow 0$ come into ∂_x and ∂_t , respectively.

We will say that a given operator $\mathcal{O}$ is a symmetry of the linear equation $E\phi(x, t) = 0$ if $\mathcal{O}$ transforms solutions into solutions, that is, if $\mathcal{O}$ is such that

$$E\mathcal{O} = \Lambda E, \quad (3)$$

where Λ is another operator. In this way, the symmetries of the equation (1) were computed,⁹ showing that they realized the Schrödinger algebra $\mathcal{S}$, which is exactly the same result as for the continuous case.^{17,18}

On the other hand, two quantum deformations of the Schrödinger algebra have been recently obtained,^{19,20} both of them endowed with a triangular Hopf structure. The former was derived by starting from the non-standard quantum deformation of its harmonic oscillator subalgebra h_4 , while the latter was constructed by means of the non-standard quantum deformation of the extended $\overline{sl}(2, \mathbb{R})$ subalgebra. In Sec. II we show that the symmetry algebra⁹ of the space discretization of the SE obtained from (1) by taking the limit $\tau \rightarrow 0$ is just the quantum Schrödinger algebra given by Ballesteros *et al.*,¹⁹ and the deformation parameter z is related with the space lattice constant σ . Likewise, we also show in Sec. III that the time discretization of the SE obtained from (1) by means of the limit $\sigma \rightarrow 0$ has the quantum Schrödinger algebra obtained by Ballesteros *et al.*²⁰ as its symmetry algebra; in this case, the time lattice step τ plays the role of the deformation parameter. As a consequence, a relationship between non-standard deformations and regular lattice discretizations can be established. In order to derive both results, the introduction of nonlinear maps transforming the aforementioned quantum algebras into non-cocommutative Hopf algebras with non-deformed commutation rules turns out to be essential. Such nonlinear transformations are explicitly given and, in the case of the quantum algebra just mentioned,²⁰ it is used to derive a new nonlinear map for the jordanian quantum deformation of $sl(2, \mathbb{R})$. Finally, the problem of finding a quantum algebra related to a full space-time discretization of the SE is discussed in Sec. IV.

II. A DISCRETE SPACE SCHRÖDINGER EQUATION

We will use the familiar notation for the Schrödinger generators: time translation H , space translation P , Galilean boost K , dilation D , conformal transformation C , and the central generator M .^{17,18} Let $U_z^{(s)}(\mathcal{S})$ be the quantum Schrödinger algebra obtained by Ballesteros *et al.*¹⁹ whose underlying Lie bialgebra is generated by the non-standard classical r -matrix

$$r^{(s)} = zP \wedge (D + \frac{1}{2}M). \quad (4)$$

The coproduct of $U_z^{(s)}(\mathcal{S})$ is

$$\begin{aligned} \Delta(M) &= 1 \otimes M + M \otimes 1 \\ \Delta(P) &= 1 \otimes P + P \otimes 1 \\ \Delta(H) &= 1 \otimes H + H \otimes e^{-2zP} \\ \Delta(K) &= 1 \otimes K + K \otimes e^{zP} - zD' \otimes e^{zP}M \\ \Delta(D) &= 1 \otimes D + D \otimes e^{zP} + \frac{1}{2}M \otimes (e^{zP} - 1) \\ \Delta(C) &= 1 \otimes C + C \otimes e^{2zP} + \frac{z}{2}K \otimes e^{zP}D' - \frac{z}{2}D' \otimes e^{zP}(K + zD'M), \end{aligned} \quad (5)$$

and its commutation rules are

$$\begin{aligned}
[D, P] &= \frac{1}{z}(1 - e^{zP}) & [D, K] &= K & [K, P] &= Me^{zP} & [M, \cdot] &= 0 \\
[D, H] &= -2H & [D, C] &= 2C - \frac{z}{2}KD' & [H, P] &= 0 \\
[H, C] &= \frac{1}{2}(1 + e^{-zP})D' - \frac{1}{2}M + zKH & [K, C] &= -\frac{z}{2}K^2 \\
[P, C] &= -\frac{1}{2}(1 + e^{zP})K - \frac{z}{2}e^{zP}MD' & [K, H] &= \frac{1}{z}(1 - e^{-zP})
\end{aligned} \tag{6}$$

where hereafter we shall use the notation $D' = D + \frac{1}{2}M$ in order to simplify some expressions. The following differential-difference realization of (6) can be found:

$$\begin{aligned}
P &= \partial_x & H &= \partial_t & M &= m \\
K &= -t \frac{1 - e^{-z\partial_x}}{z} - mxe^{z\partial_x} & D &= 2t\partial_t + x \frac{e^{z\partial_x} - 1}{z} + \frac{1}{2} \\
C &= t^2\partial_t e^{-z\partial_x} + tx \left(\frac{\sinh z\partial_x}{z} - zm\partial_t e^{z\partial_x} \right) + \frac{1}{2}mx^2 e^{z\partial_x} \\
&\quad - \frac{1}{4}t \left\{ 1 - 3e^{-z\partial_x} + m(1 - e^{-z\partial_x}) \right\} + \frac{1}{4}zm(1 - m)xe^{z\partial_x}.
\end{aligned} \tag{7}$$

Note that the Galilei generators $\{K, H, P, M\}$ close a deformed subalgebra (but not a Hopf subalgebra) whose Casimir is

$$E_z^{(s)} = \left(\frac{1 - e^{-zP}}{z} \right)^2 - 2MH. \tag{8}$$

The action of $E_z^{(s)}$ on $\phi(x, t)$ through (7) provides a space discretization of the SE:

$$\left(\left(\frac{1 - e^{-z\partial_x}}{z} \right)^2 - 2m\partial_t \right) \phi(x, t) = 0. \tag{9}$$

Furthermore, the quantum Hopf algebra $U_z^{(s)}(\mathcal{S})$ is a symmetry algebra of this equation since

$$\begin{aligned}
[E_z^{(s)}, X] &= 0 \quad \text{for } X \in \{K, H, P, M\} & [E_z^{(s)}, D] &= 2E_z^{(s)} \\
[E_z^{(s)}, C] &= \left\{ t(e^{-z\partial_x} + 1) - zmx e^{z\partial_x} \right\} E_z^{(s)}.
\end{aligned} \tag{10}$$

Now we will relate these results with the discretizations of the SE studied by Floreanini *et al.*⁹ through the non-linear change of basis defined by

$$\begin{aligned}
\mathcal{H} &= H & \mathcal{P} &= \frac{1 - e^{-zP}}{z} & \mathcal{M} &= M \\
\mathcal{D} &= D + \frac{1}{2}(1 - e^{zP}) & \mathcal{K} &= -2K - zMe^{zP} \\
\mathcal{C} &= C - \frac{z}{2}KD' + \frac{z}{2}Ke^{zP} - \frac{z^2}{8}Me^{2zP} & \sigma &= -z.
\end{aligned} \tag{11}$$

In this new basis the commutation rules (6) of the Hopf algebra $U_\sigma^{(s)}(\mathcal{S})$ are

$$\begin{aligned}
[\mathcal{D}, \mathcal{P}] &= -\mathcal{P} & [\mathcal{D}, \mathcal{K}] &= \mathcal{K} & [\mathcal{K}, \mathcal{P}] &= -2\mathcal{M} & [\mathcal{M}, \cdot] &= 0 \\
[\mathcal{D}, \mathcal{H}] &= -2\mathcal{H} & [\mathcal{D}, \mathcal{C}] &= 2\mathcal{C} & [\mathcal{H}, \mathcal{C}] &= \mathcal{D} & [\mathcal{H}, \mathcal{P}] &= 0 \\
[\mathcal{P}, \mathcal{C}] &= \frac{1}{2}\mathcal{K} & [\mathcal{K}, \mathcal{H}] &= -2\mathcal{P} & [\mathcal{K}, \mathcal{C}] &= 0
\end{aligned} \tag{12}$$

that is, they come into the non-deformed Schrödinger Lie algebra. Hence the deformation parameter σ appears only within the coproduct, which now reads

$$\begin{aligned}
\Delta(\mathcal{M}) &= 1 \otimes \mathcal{M} + \mathcal{M} \otimes 1 \\
\Delta(\mathcal{P}) &= 1 \otimes \mathcal{P} + \mathcal{P} \otimes 1 + \sigma \mathcal{P} \otimes \mathcal{P} \\
\Delta(\mathcal{H}) &= 1 \otimes \mathcal{H} + \mathcal{H} \otimes (1 + \sigma \mathcal{P})^2 \\
\Delta(\mathcal{K}) &= 1 \otimes \mathcal{K} + \mathcal{K} \otimes \frac{1}{1 + \sigma \mathcal{P}} - 2\sigma \mathcal{D}' \otimes \frac{\mathcal{M}}{1 + \sigma \mathcal{P}} \\
\Delta(\mathcal{D}) &= 1 \otimes \mathcal{D} + \mathcal{D} \otimes \frac{1}{1 + \sigma \mathcal{P}} - \frac{1}{2} \mathcal{M} \otimes \frac{\sigma \mathcal{P}}{1 + \sigma \mathcal{P}} \\
\Delta(\mathcal{C}) &= 1 \otimes \mathcal{C} + \mathcal{C} \otimes \frac{1}{(1 + \sigma \mathcal{P})^2} - \frac{\sigma}{2} \mathcal{D}' \otimes \frac{1}{1 + \sigma \mathcal{P}} \mathcal{K} \\
&\quad + \frac{\sigma^2}{2} \mathcal{D}'(\mathcal{D}' - 1) \otimes \frac{\mathcal{M}}{(1 + \sigma \mathcal{P})^2}
\end{aligned} \tag{13}$$

where $\mathcal{D}' = \mathcal{D} + \frac{1}{2}\mathcal{M}$. Note that although the new generator $\mathcal{P}$ is non-primitive, its coproduct satisfies

$$\Delta((1 + \sigma \mathcal{P})^a) = (1 + \sigma \mathcal{P})^a \otimes (1 + \sigma \mathcal{P})^a \tag{14}$$

since the old generator P fulfils $\Delta(e^{azP}) = e^{azP} \otimes e^{azP}$ for any real number a .

On the other hand, the mapping (11) transforms the differential-difference realization (7) into

$$\begin{aligned}
\mathcal{P} &= \Delta_x & \mathcal{H} &= \partial_t & \mathcal{M} &= m \\
\mathcal{K} &= 2t\Delta_x + m(2x + \sigma)T_x^{-1} & \mathcal{D} &= 2t\partial_t + x\Delta_x T_x^{-1} - \frac{1}{2}T_x^{-1} + 1 \\
\mathcal{C} &= t^2\partial_t + tx\Delta_x T_x^{-1} + \frac{1}{2}mx^2 T_x^{-2} + t\left(1 - \frac{1}{2}T_x^{-1}\right) - \frac{m\sigma^2}{8}T_x^{-2}.
\end{aligned} \tag{15}$$

We emphasize that these are just the symmetry operators of the equation (1) obtained by Floreanini *et al.*⁹, provided the continuum time limit $\tau \rightarrow 0$ is performed and $m = \frac{1}{2}$. This connection is consistent with the fact that (11) transforms the Casimir of the Galilei subalgebra $E_z^{(s)}$ (8) into the non-deformed one

$$E = \mathcal{P}^2 - 2\mathcal{M}\mathcal{H}. \tag{16}$$

Therefore, the discretized SE

$$(\Delta_x^2 - 2m\partial_t)\phi(x, t) = 0 \tag{17}$$

is obtained as the realization (15) of E , and the operators (15) are symmetries of this equation satisfying

$$[E, X] = 0 \quad X \in \{\mathcal{K}, \mathcal{H}, \mathcal{P}, \mathcal{M}\} \quad [E, \mathcal{D}] = 2E \quad [E, \mathcal{C}] = 2tE. \tag{18}$$

Thus, we have shown that the space differential-difference SE under study⁹ has $U_\sigma^{(s)}(\mathcal{S})$ as its quantum Hopf symmetry algebra. The deformation parameter $\sigma = -z$ is interpreted as the lattice step in the x coordinate, meanwhile the time t remains as a continuum variable. We also remark that, by using (15), the solutions of (17) have been obtained⁹ for $m = \frac{1}{2}$.

III. A DISCRETE TIME SCHRÖDINGER EQUATION

A similar procedure can be applied to the quantum Schrödinger algebra $U_z^{(t)}(\mathcal{S})$ coming from the non-standard classical r -matrix

$$r^{(t)} = 2zH \wedge (D + \frac{1}{2}M) \equiv 2zH \wedge D'. \quad (19)$$

The coproduct and commutation rules of $U_z^{(t)}(\mathcal{S})$ are given by²⁰

$$\begin{aligned} \Delta(M) &= 1 \otimes M + M \otimes 1 \\ \Delta(H) &= 1 \otimes H + H \otimes 1 \\ \Delta(P) &= 1 \otimes P + P \otimes e^{-2zH} \\ \Delta(K) &= 1 \otimes K + K \otimes e^{2zH} - 2zD' \otimes e^{4zH}P \\ \Delta(D) &= 1 \otimes D + D \otimes e^{4zH} + \frac{1}{2}M \otimes (e^{4zH} - 1) \\ \Delta(C) &= 1 \otimes C + C \otimes e^{4zH} - zD' \otimes e^{4zH}M \end{aligned} \quad (20)$$

$$\begin{aligned} [D, P] &= -P & [D, K] &= K & [K, P] &= M & [M, \cdot] &= 0 \\ [D, H] &= \frac{1}{2z}(1 - e^{4zH}) & [D, C] &= 2C + 2z(D')^2 & [H, P] &= 0 \\ [H, C] &= D' - \frac{1}{2}Me^{4zH} & [K, C] &= z(D'K + KD') \\ [P, C] &= -K - z(D'P + PD') & [K, H] &= e^{4zH}P. \end{aligned} \quad (21)$$

A realization of (21) reads²⁰

$$\begin{aligned} H &= \partial_t & P &= \partial_x & M &= m \\ K &= -(t + 4z)e^{4z\partial_t}\partial_x - mx & D &= 2(t + 4z)\frac{e^{4z\partial_t} - 1}{4z} + x\partial_x + \frac{1}{2} \\ C &= (t^2 - 4zbt)\frac{e^{4z\partial_t} - 1}{4z} + tx\partial_x + \frac{1}{2}t + \frac{1}{2}mx^2 - 4z(b + 1)e^{4z\partial_t} \\ &\quad - zx^2\partial_x^2 - 2z(b + 1)x\partial_x - z(b + 1/2)^2, \end{aligned} \quad (22)$$

where $b = \frac{m}{2} - 2$. Now, a time discretization of the SE is obtained by considering the deformed Casimir of the Galilei subalgebra

$$E_z^{(t)} = P^2 - 2M \left(\frac{1 - e^{-4zH}}{4z} \right) \quad (23)$$

written in terms of the realization (22):

$$\left(\partial_x^2 - 2m \frac{1 - e^{-4z\partial_t}}{4z} \right) \phi(x, t) = 0. \quad (24)$$

Under the realization (22) the generators of $U_z^{(t)}(\mathcal{S})$ are symmetry operators of this equation as they satisfy

$$\begin{aligned} [E_z^{(t)}, X] &= 0 \quad X \in \{K, H, P, M\} & [E_z^{(t)}, D] &= 2E_z^{(t)} \\ [E_z^{(t)}, C] &= 2\{t + z(1 - m - 2x\partial_x)\} E_z^{(t)}. \end{aligned} \quad (25)$$

The relationship with the time discretization of the SE analysed by Floreanini *et al.*⁹ is provided by the non-linear map defined by

$$\begin{aligned} \mathcal{H} &= \frac{1 - e^{-4zH}}{4z} & \mathcal{P} &= P & \mathcal{M} &= M \\ \mathcal{D} &= D + 2(1 - e^{4zH}) & \mathcal{K} &= -2K - 8zPe^{4zH} \\ \mathcal{C} &= C + z(D')^2 - 4zDe^{4zH} & \tau &= -4z. \end{aligned} \quad (26)$$

If we apply (26) to (21) we find again the classical commutation rules of the Schrödinger algebra (12) while the coproduct is now given by

$$\begin{aligned} \Delta(\mathcal{M}) &= 1 \otimes \mathcal{M} + \mathcal{M} \otimes 1 \\ \Delta(\mathcal{H}) &= 1 \otimes \mathcal{H} + \mathcal{H} \otimes 1 + \tau \mathcal{H} \otimes \mathcal{H} \\ \Delta(\mathcal{P}) &= 1 \otimes \mathcal{P} + \mathcal{P} \otimes (1 + \tau \mathcal{H})^{1/2} \\ \Delta(\mathcal{K}) &= 1 \otimes \mathcal{K} + \mathcal{K} \otimes \frac{1}{(1 + \tau \mathcal{H})^{1/2}} - \tau \mathcal{D}' \otimes \frac{\mathcal{P}}{1 + \tau \mathcal{H}} \\ \Delta(\mathcal{D}) &= 1 \otimes \mathcal{D} + \mathcal{D} \otimes \frac{1}{1 + \tau \mathcal{H}} - \frac{1}{2} \mathcal{M} \otimes \frac{\tau \mathcal{H}}{1 + \tau \mathcal{H}} \\ \Delta(\mathcal{C}) &= 1 \otimes \mathcal{C} + \mathcal{C} \otimes \frac{1}{1 + \tau \mathcal{H}} - \frac{\tau}{2} \mathcal{D}' \otimes \frac{1}{1 + \tau \mathcal{H}} \mathcal{D} \\ &\quad + \frac{\tau}{4} \mathcal{D}'(\mathcal{D}' - 2) \otimes \frac{\tau \mathcal{H}}{(1 + \tau \mathcal{H})^2}. \end{aligned} \quad (27)$$

The generator $\mathcal{H}$ fulfils a similar property to (14), that is,

$$\Delta((1 + \tau \mathcal{H})^a) = (1 + \tau \mathcal{H})^a \otimes (1 + \tau \mathcal{H})^a. \quad (28)$$

On the other hand, in this new basis the realization (22) turns out to be

$$\begin{aligned} \mathcal{H} &= \Delta_t & \mathcal{P} &= \partial_x & \mathcal{M} &= m \\ \mathcal{K} &= 2t\partial_x T_t^{-1} + 2mx & \mathcal{D} &= 2t\Delta_t T_t^{-1} + x\partial_x + \frac{1}{2} \\ \mathcal{C} &= t^2 \Delta_t T_t^{-2} + tx\partial_x T_t^{-1} + \frac{1}{2}mx^2 + t\left(T_t^{-2} - \frac{1}{2}T_t^{-1}\right), \end{aligned} \quad (29)$$

which is again the limit $\sigma \rightarrow 0$ of the symmetry operators of Floreanini *et al.*⁹ when $m = \frac{1}{2}$. The corresponding discretized SE is provided by the (non-deformed) Casimir E of the Galilei subalgebra (16) written through (29):

$$(\partial_x^2 - 2m\Delta_t)\phi(x, t) = 0. \quad (30)$$

The new operators (29) are symmetries of this equation satisfying

$$[E, X] = 0 \quad X \in \{\mathcal{K}, \mathcal{H}, \mathcal{P}, \mathcal{M}\} \quad [E, \mathcal{D}] = 2E \quad [E, \mathcal{C}] = 2tT_t^{-1}E. \quad (31)$$

Thus we have obtained a discrete version of the SE on a uniform lattice such that its symmetry algebra is the quantum Hopf algebra $U_\tau^{(t)}(\mathcal{S})$, and the time lattice step is related to the deformation parameter in the form $\tau = -4z$.

A. A NONLINEAR MAP FOR THE JORDANIAN DEFORMATION OF $sl(2, \mathbb{R})$

We emphasize that the four generators $\{\mathcal{D}, \mathcal{H}, \mathcal{C}, \mathcal{M}\}$ close a quantum extended $sl(2, \mathbb{R})$ algebra. Hence, as a byproduct, if we take $\mathcal{M} = 0$ we shall get the map

$$\begin{aligned} \mathcal{H} &= \frac{1 - e^{-4zH}}{4z} \\ \mathcal{D} &= D + 2(1 - e^{4zH}) \\ \mathcal{C} &= C + zD^2 - 4zDe^{4zH} \quad \tau = -4z. \end{aligned} \quad (32)$$

Now, if we denote

$$J_3 = -\mathcal{D} \quad J_+ = \mathcal{H} \quad J_- = -\mathcal{C}, \quad (33)$$

and we apply this map onto the well known non-standard quantum $sl(2, \mathbb{R})$ algebra^{11–16} given from (20) and (21)

$$\begin{aligned} \Delta(H) &= 1 \otimes H + H \otimes 1 \\ \Delta(D) &= 1 \otimes D + D \otimes e^{4zH} \\ \Delta(C) &= 1 \otimes C + C \otimes e^{4zH} \end{aligned} \quad (34)$$

$$[D, H] = \frac{1}{2z}(1 - e^{4zH}) \quad [D, C] = 2C + 2z(D)^2 \quad [H, C] = D, \quad (35)$$

we obtain a new expression for this quantum algebra with deformed non-cocommutative coproduct and “classical” commutation rules:

$$\begin{aligned} [J_3, J_+] &= 2J_+ \quad [J_3, J_-] = -2J_- \quad [J_+, J_-] = J_3 \\ \Delta(J_+) &= 1 \otimes J_+ + J_+ \otimes 1 + \tau J_+ \otimes J_+ \\ \Delta(J_3) &= 1 \otimes J_3 + J_3 \otimes \frac{1}{1 + \tau J_+} \\ \Delta(J_-) &= 1 \otimes J_- + J_- \otimes \frac{1}{1 + \tau J_+} + \frac{\tau}{2} J_3 \otimes \frac{1}{1 + \tau J_+} J_3 \\ &\quad - \frac{\tau}{4} J_3 (J_3 + 2) \otimes \frac{\tau J_+}{(1 + \tau J_+)^2}, \end{aligned} \quad (36)$$

where the underlying classical r -matrix is $r = \frac{\tau}{2} J_+ \wedge J_3$. This result is worthy to be compared with the previous literature on nonlinear maps for the non-standard quantum $sl(2, \mathbb{R})$ algebra of Abdessalam *et al.*^{21,22} since, in general, the transformed coproduct has

a very complicated form (in this respect see the aforementioned references and also the work of Aizawa,²³ where the corresponding map is used to construct the representation theory of this quantum algebra).

IV. ON A POSSIBLE SPACE-TIME DISCRETIZATION

It would be certainly interesting to analyse whether there exists a (at least two parameter) quantum Schrödinger algebra giving rise to the symmetries of the full space-time uniform discretization provided by (1). In general, the search for quantum deformations of a given Lie algebra can be guided by the study of its Lie bialgebra structures, since they are just the first order deformations of the coproduct.²⁴ Therefore, in our case, one could try to find the most general Schrödinger bialgebra $(\mathcal{S}, \delta)$ such that it includes among its components the two Lie bialgebras linked to the previous deformations; hence $(\mathcal{S}, \delta)$ should depend on σ , τ and perhaps on some additional parameters λ_i . Under such conditions and by imposing δ to fulfil the cocycle condition and the dual map δ^* to define a Lie bracket, we obtain a unique cocommutator family depending only on three parameters: $\tau = -4z_1$, $\sigma = -z_2$ and λ . Furthermore, it can be shown that this Schrödinger bialgebra is a coboundary one generated by the classical r -matrix:

$$\begin{aligned}
r = & 2z_1 H \wedge D' + z_2 P \wedge D' - \lambda H \wedge C - \frac{\lambda^2}{8z_1} C \wedge D - \frac{2z_1 z_2}{\lambda} P \wedge H \\
& + \frac{\lambda z_2}{8z_1} \left(K \wedge D - \frac{\lambda}{2z_1} K \wedge C - 3P \wedge C \right) + \left(\frac{\lambda}{4} + \frac{3z_2^2}{16z_1} \right) K \wedge P \\
& + \left(\frac{3\lambda z_2}{16z_1} + \frac{z_2^3}{32z_1^2} \right) K \wedge M - \left(\frac{\lambda^2}{16z_1} + \frac{\lambda z_2^2}{32z_1^2} \right) M \wedge C \\
& + \left(\frac{\lambda}{4} + \frac{z_2^2}{16z_1} \right) M \wedge D.
\end{aligned} \tag{37}$$

We omit the explicit expressions for the cocommutators as they can be obtained by means of $\delta(X) = [1 \otimes X + X \otimes 1, r]$.

We remark that the first two terms of (37) correspond to the time and space classical r -matrices $r^{(t)}$ and $r^{(s)}$, but both deformations do not fulfil a simple “superposition principle” and extra contributions have to be added. The full quantization of such Lie bialgebra seems to be a difficult task and we shall not address this problem here. The “additional” deformation parameter λ allows us to distinguish between non-standard (when $\lambda = -\frac{z_2^2}{4z_1}$) and standard solutions (otherwise). On the other hand, although the limits $z_2 \rightarrow 0$ and $\lambda \rightarrow 0$ lead to the time classical r -matrix, unfortunately the limit $z_1 \rightarrow 0$ leads to divergences in (37).

V. CONCLUDING REMARKS

We wish to point out that the existence of a Hopf structure for the symmetries of a given equation associated to an elementary system allows us to write equations of composed systems keeping the same symmetry algebra.^{25,26} In order to use this property for the two cases here discussed, we see that only the last commutator in either (18) or (31) involving the conformal generator $\mathcal{C}$ is not algebraic, but depends explicitly on the chosen representation (the same happens at the continuum level). Consequently the composed systems characterized by the equation $\Delta(E)\phi = 0$ will have, by construction, $\Delta(\mathcal{H}), \Delta(\mathcal{P}), \Delta(\mathcal{K}), \Delta(\mathcal{D})$, and $\Delta(\mathcal{M})$ as symmetry operators (moreover they close a Hopf subalgebra). However, in general this will not be the case for $\Delta(\mathcal{C})$, and a further study on the behaviour of this operator is needed in order to construct coupled equations with full Schrödinger algebra symmetry.

Finally, we wish to point out that the applicability of the constructive approach presented here is not limited to the cases analyzed before; in fact it can be directly extended to other quantum algebras by means of their corresponding difference realizations. Work on this line is nowadays in progress.

ACKNOWLEDGEMENTS

This work has been partially supported by DGES (Projects PB94-1115 and PB95-0719) from the Ministerio de Educación y Cultura (Spain), and also by the Junta de Castilla y León (Projects CO1/396, CO2/197, and CO2/297).

¹ J. Lukierski, H. Ruegg and A. Nowicky, Phys. Lett. B **293**, 344 (1992).

² L. Frappat and A. Sciarrino, Phys. Lett. B **347**, 1 (1995).

³ A. Ballesteros, F.J. Herranz, M.A. del Olmo and M. Santander, Phys. Lett. B **351**, 137 (1995).

⁴ G.R.W. Quispel, H.W. Capel and R. Sahadevan, Phys. Lett. A **170**, 379 (1992).

⁵ D. Levi, L. Vinet and P. Winternitz, J. Phys. A **30**, 633 (1997).

⁶ R. Floreanini and L. Vinet, Lett. Math. Phys. **32**, 37 (1994).

⁷ R. Floreanini and L. Vinet, J. Math. Phys. **36**, 3134 (1995).

- ⁸ V.K. Dobrev, H.D. Doebner and C. Mrugalla, J. Phys. A **29**, 5909 (1996).
- ⁹ R. Floreanini, J. Negro, L.M. Nieto and L. Vinet, Lett. Math. Phys. **36**, 351 (1996).
- ¹⁰ J. Negro and L.M. Nieto, J. Phys. A **29**, 1107 (1996).
- ¹¹ S. Majid, Class. Quantum Grav. **5**, (1988) 1587.
- ¹² Demidov E E, Manin Yu I, Mukhin E E and Zhdanovich D V 1990 *Progr. Theor. Phys. Suppl.* **102** 203
- ¹³ C. Ohn, Lett. Math. Phys. **25**, 85 (1992).
- ¹⁴ A.A. Vladimirov, Mod. Phys. Lett. A **8**, 2573 (1993).
- ¹⁵ A. Shariati, A. Aghamohammadi and M. Khorrami, Mod. Phys. Lett. A **11**, 187 (1996).
- ¹⁶ A. Ballesteros and F.J. Herranz, J. Phys. A **29**, L311 (1996).
- ¹⁷ C.R. Hagen, Phys. Rev. D **5**, 377 (1972).
- ¹⁸ U. Niederer, Helv. Phys. Acta **45**, 802 (1972).
- ¹⁹ A. Ballesteros, F.J. Herranz and P. Parashar, J. Phys. A **30**, 8587 (1997).
- ²⁰ A. Ballesteros, F.J. Herranz and P. Parashar, Mod. Phys. Lett. A **13**, 1241 (1998).
- ²¹ B. Abdesselam, A. Chakrabarti and R. Chakrabarti, Mod. Phys. Lett. A **11**, 2883 (1996).
- ²² B. Abdesselam, A. Chakrabarti and R. Chakrabarti, Mod. Phys. Lett. A **13**, 779 (1998).
- ²³ N. Aizawa, J. Phys. A **30**, 5981 (1997).
- ²⁴ A. Ballesteros and F.J. Herranz, J. Phys. A **29**, 4307 (1996).
- ²⁵ F. Bonechi, E. Celeghini, R. Giachetti, E. Sorace and M. Tarlini, Phys Rev Lett **68**, 3718 (1992).
- ²⁶ F. Bonechi, E. Celeghini, R. Giachetti, E. Sorace and M. Tarlini, Phys Rev B **46**, 5727 (1992).